

Performing Polymerase Chain Reaction to determine the prevalence of different SAgS in *Staphylococcus aureus* samples.

Natasha Bennett, Dr. Amanda Brosnahan, Dr. Taylor Mach
Department of Science, Concordia University, St. Paul

Overview

Staphylococcus aureus is a bacteria that lives commensally on one third of the human population. *S. aureus* can produce toxins and superantigens, like SEA, TSST-1 and alpha toxin that cause disease. Some diseases produced by these toxins are pneumonia, endocarditis, osteomyelitis, toxic shock syndrome, dermatitis, and sepsis. Twelve *S. aureus* positive samples, previously collected from nasal swabs, were tested through PCR and DNA gel electrophoresis in order to visualize the DNA you are looking for.

Methods

- Twelve positive *S. aureus* samples from the study were tested for toxins
 - Isolate DNA from *S. aureus* samples
 - PCR
 - gel electrophoresis
- PCR allows the DNA of a sample to be amplified allowing it to be studied, including determining if toxins are present



Alpha toxin is present in most nasal-derived *Staphylococcus aureus* samples




Figure 1: DNA gel electrophoresis of alpha toxin on *S. aureus* samples. A DNA gel was imaged using GelGreen and blue light UV illumination to show which *S. aureus* samples produced alpha toxin. All samples presented a band at 2,000 Kb showing the presence of alpha toxin.

Background

- Alpha toxin is a pore forming toxin and when it binds to its target cell it puts a β -barrel through the target cell's membrane making a channel, causing rapid release of cellular ions.
- Alpha toxin can bind and attack a wide variety of cell causing large inflammatory response and cell death, leading to shock and sepsis.
- TSST-1 and SEA are superantigens that crosslink the V β domain of lymphocytes and class II MHC, impacting the T-cell response.
- TSST-1 and SEA causes a burst in cytokine release, also known a cytokine storm, leading to an extreme inflammatory response.
- Due to the high proinflammatory response, causing symptoms of high fever, sepsis and more.




Figure 2: DNA gel electrophoresis of the superantigen SEA on *S. aureus* samples. A DNA gel was imaged using GelGreen and blue light UV illumination to show which *S. aureus* samples produced SEA. All samples, except S0147, presented a band at 1,500 Kb showing the presence of SEA.

Results

- All 12 samples showed presence of alpha toxin
- 11:12 samples showed presence of SEA
- No samples showed presence of TSST-1

Acknowledgements & References

This research was partially funded by seven CSP Faculty Development Grants. This work has IRB approval from CSP (studies 2016_42 & 2018_37).

Pinchuk, I. V., Beswick, E. J., & Reyes, V. E. (2010). Staphylococcal enterotoxins. *Toxins*, 2(8), 2177–2197. <https://doi.org/10.3390/toxins2082177>

Oliveira, D., Borges, A., & Simões, M. (2018). *Staphylococcus aureus* Toxins and Their Molecular Activity in Infectious Diseases. *Toxins*, 10(6), 252. <https://doi.org/10.3390/toxins10060252> Staphylococcus aureus in Healthcare Settings. (2011, January 17). Retrieved April 3, 2020, from <https://www.cdc.gov/hai/organisms/staph.html>

Toxic Shock Syndrome Toxin 1. (n.d.). Retrieved April 3, 2020, from <https://www.sciencedirect.com/topics/medicine-and-dentistry/toxic-shock-syndrome-toxin-1>